

with early surgical resection of endobronchial metastasis in a follow-up of ovarian carcinoma. *Radiol Bras.* 2015;48:130.

8. Barbosa BC, Marchiori E, Zanetti GMR, et al. Catamenial pneumothorax. *Radiol Bras.* 2015;48:128–9.

9. Francisco FAF, Rodrigues RS, Barreto MM, et al. Can chest high-resolution computed tomography findings diagnose pulmonary alveolar microlithiasis? *Radiol Bras.* 2015;48:205–10.

10. Batista MN, Barreto MM, Cavaguti RF, et al. Pulmonary artery sarcoma mimicking chronic pulmonary thromboembolism. *Radiol Bras.* 2015;48:333–4.

11. Fernandes GL, Teixeira AA, Antón AGS, et al. Churg-Strauss syndrome: a case report. *Radiol Bras.* 2014;47:259–61.

12. Nishiyama KH, Falcão EAA, Kay FU, et al. Acute tracheobronchitis caused by *Aspergillus*: case report and imaging findings. *Radiol Bras.* 2014;47:317–9.

Antonio Gabriel de Jesus Barbosa¹, Diana Pena², Gláucia Zanetti¹, Edson Marchiori¹

1. Universidade Federal do Rio de Janeiro (UFRJ), Rio de Janeiro, RJ, Brazil. 2. Heart and Chest Hospital NHS Foundation Trust, Liverpool, UK. Mailing address: Dr. Edson Marchiori. Rua Thomaz Cameron, 438, Valparaíso. Petrópolis, RJ, Brazil, 25685-120. E-mail: edmarchiori@gmail.com.

<http://dx.doi.org/10.1590/0100-3984.2015.0169>

Malignant uterine disease with concurrent myometrial contraction at MRI: a possible source of overstaging

Dear Editor,

We report a case of a 41-year-old woman with lower abdominal pain and vaginal bleeding with negative ultrasound scan except for fibroids, and elevation of serum β -hCG (244,410 mIU/mL). Differential diagnoses included ectopic pregnancy, early pregnancy failure, very early ongoing pregnancy and molar pregnancy.

MRI showed an enlarged uterus with central heterogeneous T2 hyperintensity distending the endometrial canal, demonstrating reticular enhancement, concerning for gestational trophoblastic disease. There was also distortion of the junctional zone with broad intermediate-to-low signal on T2-weighted images. The

variable appearance of myometrial thickness especially on post-contrast images, showing a homogeneous myometrium, facilitated the diagnosis of contractions (Figure 1). The patient underwent suction and curettage with a final diagnosis of complete hydatidiform mole (HM) (p57 negative).

Gestational trophoblastic disease (GTD) arise from placental trophoblastic tissue after abnormal fertilization and comprises a spectrum of disorders from the pre-malignant conditions of partial HM and complete HM to the malignant invasive mole, choriocarcinoma and the very rare placental site trophoblastic tumor^(1,2). HM is the most common manifestation of GTD (85%) and by definition noninvasive and confined to the endometrium. A HM that invades the myometrium is termed invasive mole, and is composed of HM villi within the myometrium. Chorioadenoma




Figure 1. Pelvic MRI. Axial fat-suppressed (A) and sagittal (B) T2-weighted images, and post-contrast axial (C) and sagittal fat-suppressed (D) T1-weighted MRI images. An enlarged uterus is depicted, with a heterogeneous T2-weighted hyperintense lesion distending the endometrial canal (arrows, A-D). The lesion shows reticular enhancement on postcontrast imaging, concerning for gestational trophoblastic disease (short arrows, C and D). Note the different morphologic aspect of the anterior and posterior uterine walls between the first set of images (A and B) and those acquired later (C and D), suggesting motion in the context of contraction. These differences are more accentuated on sagittal images, showing substantial increase in thickness and bulging of the posterior myometrial wall on T2-weighted image (B), whereas this pattern is inverted and appearing on the anterior myometrial wall on late post-contrast T1-weighted images (D) (acquired with a delay of 30 minutes compared to T2-weighted images).

destruens is a locally invasive (myometrium) manifestation of complete HM that represents 13% of cases of GTD. Two percent of complete HM cases are described as choriocarcinoma, which is locally invasive and potentially metastasizing. These three entities produce peculiarly high levels of β -hCG, while placental site trophoblastic tumor causes a rise in human placental lactogen levels, and less elevated β -hCG levels^(3,4). Clinical assessment is difficult early in the course of the disease, as few clinical characteristics are present to distinguish it from a normal pregnancy.

Pelvic MRI is often used as a problem-solving tool in equivocal or complicated cases of GTD, especially in the first trimester, or to assess the degree of myometrial invasion and surrounding tissues^(2,5). Early manifestations appear as a soft tissue cystic mass with high T2 signal intensity⁽⁶⁾. In the second trimester these lesions tend to distend the endometrium giving a “cluster of grapes appearance”. Typically HMs are similar or slightly higher in T1 signal intensity than the adjacent myometrium. Contrast-enhanced MRI show areas of focal enhancement that relate to the amount of active trophoblastic tissue and also to β -hCG levels⁽⁷⁾. Marked early enhancement indicates active disease in the form of viable trophoblastic tissue.

In the setting of GTD, identification of myometrial invasion is crucial for diagnosis and staging. Uterine tumors associated with high serum β -hCG have a high incidence of myometrial contractions⁽⁸⁾. Myometrial contractions are seen as a bulge of the myometrial wall usually along with a region of low T2 signal intensity in the myometrium. They are transient and tend to disappear on subsequent data acquisitions⁽⁹⁾, as observed in our case. In the setting of endometrial tumor, radiologists should be aware of this phenomenon to avoid over-diagnosis and over-staging by misdiagnosing uterine contraction with myometrial extension or invasion.

REFERENCES

1. Seckl MJ, Sebire NJ, Fisher RA, et al. Gestational trophoblastic disease: ESMO Clinical Practice Guidelines for diagnosis, treatment and follow-up. *Ann Oncol.* 2013;24 Suppl 6:vi39–50.
2. Kani KK, Lee JH, Dighe M, et al. Gestational trophoblastic disease: multimodality imaging assessment with special emphasis on spectrum of abnormalities and value of imaging in staging and management of disease. *Curr Probl Diagn Radiol.* 2012;41:1–10.
3. Berkowitz RS, Goldstein DP. Diagnosis and management of the primary hydatidiform mole. *Obstet Gynecol Clin North Am.* 1988;15:491–503.
4. Ngan HY, Bender H, Benedet JL, et al. Gestational trophoblastic neoplasia, FIGO 2000 staging and classification. *Int J Gynaecol Obstet.* 2003;83 Suppl 1:175–7.
5. Allen SD, Lim AK, Seckl MJ, et al. Radiology of gestational trophoblastic neoplasia. *Clin Radiol.* 2006;61:301–13.
6. Green CL, Angtuaco TL, Shah HR, et al. Gestational trophoblastic disease: a spectrum of radiologic diagnosis. *Radiographics.* 1996;16:1371–84.
7. Yamashita Y, Torashima M, Takahashi M, et al. Contrast-enhanced dynamic MR imaging of postmolar gestational trophoblastic disease. *Acta Radiol.* 1995;36:188–92.
8. Hatakenaka M, Yoshimitsu K, Adachi T, et al. Transient uterine myometrial contraction associated with moles. *J Magn Reson Imaging.* 2004;19:182–7.
9. Ozsarlak O, Schepens E, de Schepper AM, et al. Transient uterine contraction mimicking adenomyosis on MRI. *Eur Radiol.* 1998;8:54–6.

Sriluxayini Manikkavasakar¹, Amrutha Ramachandram¹, Miguel Ramalho¹, António P. Matos¹, Richard C. Semelka¹

1. Department of Radiology, University of North Carolina at Chapel Hill, Chapel Hill, NC, USA. Mailing address: Richard C. Semelka, MD. UNC Department of Radiology. 101 Manning Drive CB#7510 – 2001 Old Clinic Bldg. Chapel Hill, NC 27599-7510, USA. E-mail: richsem@med.unc.edu.

<http://dx.doi.org/10.1590/0100-3984.2015.0057>

Leiomyoma of the breast: an uncommon tumor

Dear Editor,

A 59-year-old female patient, with no significant history, was referred by a general practitioner to our radiology clinic for routine mammography. The patient had no clinical complaints, and the physical examination revealed a painless, mobile and well-defined nodule. She underwent high-resolution mammography, which




Figure 1. In **A**, high-resolution mammogram in left craniocaudal view, and **B**, high-resolution mammogram in left mediolateral oblique view, both showing a dense nodule with lobulated contours and well-defined borders, located in the lower outer quadrant of the left breast.

identified a dense, well-defined oval nodule, located in the lower outer quadrant of the left breast (at 4 o'clock), measuring 5.5 × 3.0 cm (Figure 1). Ultrasound examination showed a well-defined oval nodule, parallel to the skin, that was hypoechoic, with no detectable Doppler flow, located in the lower outer quadrant of the left breast, measuring 3.5 × 1.7 × 3.5 cm (Figure 2). The patient underwent ultrasound-guided percutaneous core needle biopsy, and the material collected was sent for pathological study, which showed smooth muscle tumor of a benign character. In the immunohistochemical analysis, the lesion tested positive for smooth muscle actin, positive for vimentin, and negative for S100 protein, confirming the diagnosis of leiomyoma.




Figure 2. Ultrasound examination of the left breast showing a hypoechoic oval nodule with lobulated margins and well-defined borders, located in the lower outer quadrant of the left breast.